


Math 261

Fall 2023

Lecture 5



Class QZ 3

Simplify

$$\frac{\frac{1}{(x+h)^2} - \frac{1}{x^2}}{h}, \text{ evaluate your final answer with } h=0.$$

$$= \frac{x^2 \cdot 1 - (x+h)^2 \cdot 1}{x^2(x+h)^2 \cdot h}$$

$$\text{LCD} = x^2(x+h)^2$$

Box Your Final Ans


$$= \frac{x^2 - (x+h)^2}{x^2(x+h)^2 \cdot h} = \frac{x^2 - x^2 - 2xh - h^2}{x^2(x+h)^2 \cdot h} = \frac{-2xh - h^2}{x^2(x+h)^2 \cdot h}$$

$$= \frac{h(-2x - h)}{x^2(x+h)^2 \cdot h} = \frac{-2x - h}{x^2(x+h)^2}$$

for $h=0$

$$\frac{-2x - 0}{x^2(x+0)^2} = \frac{-2x}{x^2 \cdot x^2} = \frac{-2}{x^3} \checkmark$$

Intro. To Functions


Every x value can have only one y Value

$$f(x) = 3x - 2 \quad \text{Linear Function}$$

$$f(0) = 3(0) - 2 = -2 \Rightarrow (0, -2)$$

$$f(2) = 3(2) - 2 = 4 \Rightarrow (2, 4)$$

$$f(-1) = 3(-1) - 2 = -5 \Rightarrow (-1, -5)$$

as $x \rightarrow 2$ from right side $\Rightarrow f(x) = y \rightarrow 4$ as $x \rightarrow 2$ " left side $\Rightarrow f(x) = y \rightarrow 4$

$$f(x) = \sqrt{x}$$




$$x \geq 0$$

$$y \geq 0$$

Domain $[0, \infty)$ Range $[0, \infty)$

x	y
0	0
1	1
4	2
9	3

as $x \rightarrow 4$ from right, $y \rightarrow 2$ as $x \rightarrow 4^+$, $y \rightarrow 2$ as $x \rightarrow 4$

" left

, $y \rightarrow 2$ $\Rightarrow$ as $x \rightarrow 4^-$, $y \rightarrow 2$

$$f(x) = \frac{1}{x-1}$$


$$x-1 \neq 0 \rightarrow x \neq 1$$

$$\text{Domain: } (-\infty, 1) \cup (1, \infty)$$

$$f(0) = \frac{1}{0-1} = -1$$

$$f(2) = \frac{1}{2-1} = 1$$

$$\text{Range: } (-\infty, 0) \cup (0, \infty)$$



$$\text{as } x \rightarrow 1^+, y \rightarrow \infty$$


$$\text{as } x \rightarrow 1^-, y \rightarrow -\infty$$

Piece-wise Function


$$f(x) = \begin{cases} x^2 & \text{if } x < 2 \\ -4 & \text{if } x \geq 2 \end{cases}$$

$$\text{as } x \rightarrow 2^+, f(x) = -4$$

$$\text{as } x \rightarrow 2^-, f(x) \rightarrow 4$$



$$f(x) = \begin{cases} -x & \text{if } x < 1 \\ 4 & x = 1 \checkmark \\ x^2 - 1 & \text{if } x > 1 \end{cases}$$



as $x \rightarrow 1^+$, $f(x) \rightarrow 0$

as $x \rightarrow 1^-$, $f(x) \rightarrow -1$

$$f(1) = 4$$

Consider $f(x) = \frac{x^3 + 2x^2}{x+2}$


Simplify Yes

$$x+2 \neq 0 \rightarrow x \neq -2$$

modify

$$\text{Domain } (-\infty, -2) \cup (-2, \infty)$$

NO



$$f(-1.99) = \frac{(-1.99)^3 + 2(-1.99)^2}{-1.99 + 2} = 3.9601 \rightarrow 4$$

$$f(-1.999) = \frac{(-1.999)^3 + 2(-1.999)^2}{-1.999 + 2} = 3.996001 \rightarrow 4$$

$$f(-2.0001) = \frac{(-2.0001)^3 + 2(-2.0001)^2}{-2.0001 + 2} = 4.0004 \rightarrow 4$$

$$f(x) = x^2$$

Simplify

$$\frac{f(x+h) - f(x)}{h}$$

, then evaluate for $h=0$

Difference quotient

$$\begin{aligned} \frac{(x+h)^2 - x^2}{h} &= \frac{x^2 + 2xh + h^2 - x^2}{h} = \frac{2xh + h^2}{h} \\ &= \frac{\cancel{h}(2x+h)}{\cancel{h}} \\ &= 2x + h \\ &\quad \boxed{2x} \end{aligned}$$

$$f(x) = \sqrt{x}$$

Simplify the difference quotient then evaluate for

$h=0$.

$$\frac{f(x+h) - f(x)}{h}$$

$$= \frac{\sqrt{x+h} - \sqrt{x}}{h}$$

for $h=0$
 $\frac{\sqrt{x} - \sqrt{x}}{0} = \frac{0}{0}$

$$= \frac{(\sqrt{x+h} - \sqrt{x})(\sqrt{x+h} + \sqrt{x})}{h(\sqrt{x+h} + \sqrt{x})}$$

$$= \frac{\cancel{x+h} - \cancel{x}}{h(\sqrt{x+h} + \sqrt{x})}$$

Now evaluate for $h=0$

$$\begin{aligned} &= \frac{1}{\sqrt{x+0} + \sqrt{x}} = \frac{1}{\sqrt{x} + \sqrt{x}} \\ &= \boxed{\frac{1}{2\sqrt{x}}} \end{aligned}$$

$$= \frac{1}{\sqrt{x+h} + \sqrt{x}}$$

Class QZ 4

Consider $f(x) = \frac{1}{x}$

Simplify the difference quotient, then evaluate

for $h=0$. Box Your Final Answer.

$$\frac{f(x+h) - f(x)}{h} = \frac{\frac{1}{x+h} - \frac{1}{x}}{h} = \frac{x - (x+h)}{h x(x+h)} = \frac{x - x - h}{h x(x+h)}$$

$$= \frac{-h}{\cancel{h} x(x+h)} = \boxed{\frac{-1}{x(x+h)}} \quad \text{LCM} = x(x+h) \quad \text{for } h=0 \quad \frac{-1}{x(x+0)} = \frac{-1}{x \cdot x} = \boxed{\frac{-1}{x^2}}$$